

# STAAR CHEMISTRY REFERENCE MATERIALS



## ATOMIC STRUCTURE

$$\text{Speed of light} = (\text{frequency})(\text{wavelength})$$

$$c = f\lambda$$

$$\text{Energy} = (\text{Planck's constant})(\text{frequency})$$

$$E_{\text{photon}} = hf$$

$$\text{Energy} = \frac{(\text{Planck's constant})(\text{speed of light})}{(\text{wavelength})}$$

$$E_{\text{photon}} = \frac{hc}{\lambda}$$

## BEHAVIOR OF GASES

$$\text{Total pressure of a gas} = \left( \begin{array}{l} \text{sum of the partial pressures} \\ \text{of the component gases} \end{array} \right)$$

$$P_T = P_1 + P_2 + P_3 + \dots$$

$$(\text{Pressure})(\text{volume}) = (\text{moles})(\text{ideal gas constant})(\text{temperature})$$

$$PV = nRT$$

$$\frac{(\text{Initial pressure})(\text{initial volume})}{(\text{Initial moles})(\text{initial temperature})} = \frac{(\text{final pressure})(\text{final volume})}{(\text{final moles})(\text{final temperature})}$$

$$\frac{PV_1}{n_1T_1} = \frac{PV_2}{n_2T_2}$$

$$(\text{Initial pressure})(\text{initial volume}) = (\text{final pressure})(\text{final volume})$$

$$P_1V_1 = P_2V_2$$

$$\frac{(\text{Initial volume})}{(\text{Initial temperature})} = \frac{(\text{final volume})}{(\text{final temperature})}$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$\frac{(\text{Initial volume})}{(\text{Initial moles})} = \frac{(\text{final volume})}{(\text{final moles})}$$

$$\frac{V_1}{n_1} = \frac{V_2}{n_2}$$

## SOLUTIONS

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liter of solution}}$$

$$M = \frac{\text{mol}}{\text{L}}$$

$$\text{Ionization constant of water} = \left( \begin{array}{l} \text{hydrogen ion} \\ \text{concentration} \end{array} \right) \left( \begin{array}{l} \text{hydroxide ion} \\ \text{concentration} \end{array} \right)$$

$$K_w = [\text{H}^+][\text{OH}^-]$$

$$\left( \begin{array}{l} \text{Volume of} \\ \text{solution 1} \end{array} \right) \left( \begin{array}{l} \text{molarity of} \\ \text{solution 1} \end{array} \right) = \left( \begin{array}{l} \text{volume of} \\ \text{solution 2} \end{array} \right) \left( \begin{array}{l} \text{molarity of} \\ \text{solution 2} \end{array} \right)$$

$$V_1M_1 = V_2M_2$$

$$\text{pH} = -\text{log}(\text{hydrogen ion concentration})$$

$$\text{pH} = -\log[\text{H}^+]$$

## THERMOCHEMISTRY

$$\text{Heat gained or lost} = (\text{mass}) \left( \begin{array}{l} \text{specific} \\ \text{heat} \end{array} \right) \left( \begin{array}{l} \text{change in} \\ \text{temperature} \end{array} \right)$$

$$Q = mc_p \Delta T$$

$$\text{Enthalpy of reaction} = \left( \begin{array}{l} \text{enthalpy} \\ \text{of products} \end{array} \right) - \left( \begin{array}{l} \text{enthalpy} \\ \text{of reactants} \end{array} \right)$$

$$\Delta H = \Delta H_f^\circ(\text{products}) - \Delta H_f^\circ(\text{reactants})$$

# STAAR CHEMISTRY REFERENCE MATERIALS



## OTHER FORMULAS

$$\text{Density} = \frac{\text{mass}}{\text{volume}}$$

$$D = \frac{m}{V}$$

$$\text{Percent error} = \left( \frac{\text{accepted value} - \text{experimental value}}{\text{accepted value}} \right) (100)$$

$$\text{Percent yield} = \left( \frac{\text{actual yield}}{\text{theoretical yield}} \right) (100)$$

## CONSTANTS AND CONVERSIONS

$$\text{Avogadro's number} = 6.02 \times 10^{23} \text{ particles per mole}$$

$$h = \text{Planck's constant} = 6.63 \times 10^{-34} \text{ J} \cdot \text{s}$$

$$c = \text{speed of light} = 3.00 \times 10^8 \frac{\text{m}}{\text{s}}$$

$$K_w = \text{ionization constant of water} = 1.00 \times 10^{-14} \left( \frac{\text{mol}}{\text{L}} \right)^2$$

$$\text{alpha particle } (\alpha) = {}_2^4\text{He} \quad \text{beta particle } (\beta) = {}_{-1}^0\text{e} \quad \text{neutron} = {}_0^1\text{n}$$

$$\text{standard temperature and pressure (STP)} = 0^\circ\text{C and 1 atm}$$

$$0^\circ\text{C} = 273 \text{ K}$$

$$\text{volume of ideal gas at STP} = 22.4 \frac{\text{L}}{\text{mol}}$$

$$1 \text{ cm}^3 = 1 \text{ mL} = 1 \text{ cc}$$

$$1 \text{ atm} = 760 \text{ mm Hg} = 101.3 \text{ kPa}$$

$$R = \text{ideal gas constant} = 0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} = 8.31 \frac{\text{L} \cdot \text{kPa}}{\text{mol} \cdot \text{K}} = 62.4 \frac{\text{L} \cdot \text{mm Hg}}{\text{mol} \cdot \text{K}}$$

$$1 \text{ calorie (cal)} = 4.18 \text{ joules (J)}$$

$$1000 \text{ calories (cal)} = 1 \text{ Calorie (Cal)} = 1 \text{ kilocalorie (kcal)}$$

## RULES FOR SIGNIFICANT FIGURES

1. Non-zero digits and zeros between non-zero digits are always significant.
2. Leading zeros are not significant.
3. Zeros to the right of all non-zero digits are only significant if a decimal point is shown.
4. For values written in scientific notation, the digits in the coefficient are significant.
5. In a common logarithm, there are as many digits after the decimal point as there are significant figures in the original number.

# STAAR CHEMISTRY REFERENCE MATERIALS



POLYATOMIC IONS		SOLUBILITY OF COMMON IONIC COMPOUNDS IN WATER		ACTIVITY SERIES	
<u>Soluble</u> <u>compounds contain</u>		<u>Common exceptions</u>		<u>Increasing Activity</u>	
Acetate	$\text{C}_2\text{H}_3\text{O}_2^-$ , $\text{CH}_3\text{COO}^-$	$\text{C}_2\text{H}_3\text{O}_2^-$ , $\text{CH}_3\text{COO}^-$	None	Lithium	
Ammonium	$\text{NH}_4^+$	$\text{NH}_4^+$	None	Potassium	
Carbonate	$\text{CO}_3^{2-}$	$\text{NO}_3^-$	None	Barium	
Chlorate	$\text{ClO}_3^-$	$\text{CN}^-$	None	Calcium	
Chlorite	$\text{ClO}_2^-$	$\text{ClO}^-$	None	Sodium	
Chromate	$\text{CrO}_4^{2-}$	$\text{ClO}_2^-$	None	Magnesium	
Cyanide	$\text{CN}^-$	$\text{ClO}_3^-$	None	Aluminum	
Dichromate	$\text{Cr}_2\text{O}_7^{2-}$	$\text{ClO}_4^-$	None	Manganese	
Hydrogen carbonate	$\text{HCO}_3^-$	$\text{Br}^-$	Compounds of $\text{Ag}^+$ , $\text{Pb}^{2+}$ , and $\text{Hg}_2^{2+}$	Zinc	
Hydroxide	$\text{OH}^-$	$\text{Cl}^-$	Compounds of $\text{Ag}^+$ , $\text{Pb}^{2+}$ , and $\text{Hg}_2^{2+}$	Chromium	
Hypochlorite	$\text{ClO}^-$	$\text{I}^-$	Compounds of $\text{Sr}^{2+}$ , $\text{Ba}^{2+}$ , $\text{Pb}^{2+}$ , and $\text{Hg}_2^{2+}$	Iron	
Nitrate	$\text{NO}_3^-$	$\text{SO}_4^{2-}$	Compounds of $\text{Sr}^{2+}$ , $\text{Ba}^{2+}$ , $\text{Pb}^{2+}$ , and $\text{Hg}_2^{2+}$	Cobalt	
Nitrite	$\text{NO}_2^-$			Nickel	
Perchlorate	$\text{ClO}_4^-$			Tin	
Permanganate	$\text{MnO}_4^-$	$\text{Cr}_2\text{O}_7^{2-}$	Compounds of $\text{NH}_4^+$ and the alkali metal cations	Lead	
Phosphate	$\text{PO}_4^{3-}$	$\text{OH}^-$	Compounds of $\text{NH}_4^+$ and the alkali metal cations	(Hydrogen)	
Sulfate	$\text{SO}_4^{2-}$	$\text{S}^{2-}$	Compounds of $\text{NH}_4^+$ and the alkali metal cations,	Copper	
Sulfite	$\text{SO}_3^{2-}$		$\text{Ca}^{2+}$ , $\text{Sr}^{2+}$ , and $\text{Ba}^{2+}$	Mercury	
			Compounds of $\text{NH}_4^+$ , the alkali metal cations,	Silver	
			$\text{Ca}^{2+}$ , $\text{Sr}^{2+}$ , and $\text{Ba}^{2+}$	Platinum	
				Gold	

# **STAAR CHEMISTRY REFERENCE MATERIALS**



## PERIODIC TABLE OF THE ELEMENTS